2016 WCAW References

6.7 million patients are suffering with non-healing advanced wounds
(Expected to grow at 2% per year over the next decade.)

- L. E. K. Consulting Market Analysis "Market Sizing and Assessment of Outsourced Outpatient Wound Care"

There Are Approximately 29,100,000 diabetic patients in the US. 9.3% of the population.

- June 2014 American Diabetes Association Fact Sheet
- 2013 US Census bureau
- 2013 UnitedHealth Group

Each year 2.5 million patients aged 75+ acquire a pressure ulcer. 13.2% of the relevant population.

- 2011 Agency for healthcare Research and quality
- 2011 US Census bureau

Where are advanced wounds treated?
Hospital outpatient wound care: 40%
Hospital inpatient: 20%
Long term care facility: 5%
Stand along (Dr. office, ASC, home): 35%

- Healogics Management Assumption

Cost to treat advanced wounds exceeds $50 billion annually in the US

- WOUNDS 2012; 24(1): 10-17 Wound Care Outcomes and Associated Cost Among Patients Treated in US Outpatient Wound Centers; Data from the US Wound

Fact Line: Diabetic foot ulcer patients: 2,098,750; Venous Leg Ulcer patients: 817,244; pressure ulcer patients: 2,906,336; surgical/trauma wound patients: 519,923; arterial ulcer patients: 388,730

- I. E. K. Consulting Market Analysis "Market Sizing and Assessment of Outsourced Outpatient Wound Care"

30% of untreated wounds result in amputation

- Prim Care Companion J Clin Psychiatry. 2007; 9(4): 303-308 "Reactions to Amputation: Recognition and Treatment"

Patients with an amputation have a 50% morality rate within 5 years.

- Prim Care Companion J Clin Psychiatry. 2007; 9(4): 303-308 "Reactions to Amputation: Recognition and Treatment"